

Clinical Prediction for Success of Interventions for Managing Low Back Pain

Jeffrey Hebert, DC^{a,*}, Shane Koppenhaver, MPT^a,
Julie Fritz, PhD, PT, ATC^a, Eric Parent, PhD, PT^b

^aThe University of Utah, College of Health, 520 Wakara Way, Salt Lake City, UT 84108, USA

^bDepartment of Physical Therapy, The University of Alberta, Faculty of Rehabilitation Medicine, Edmonton, Canada

Low back pain (LBP) is highly prevalent in athletic and nonathletic populations, and is a common cause of pain and disability. It is difficult to identify the pathoanatomical cause for most cases of LBP, leading many to consider LBP as a single “nonspecific” disorder. Most studies evaluating the treatment effectiveness of interventions for LBP have been based on this presumption and have generally demonstrated small to no treatment effects. Most providers think of LBP as a more heterogeneous disorder, and the inability to more specifically match patients to interventions likely to be beneficial is one possible explanation for the lack of research evidence proving the effectiveness of treatments and the suboptimal outcomes of clinical care. Treatment-based classification, one approach to subgrouping patients with “nonspecific” LBP, focuses on identifying clusters of findings from the history and clinical examination that predict a more favorable outcome with a specific treatment approach. By matching patients with the appropriate specific exercise, stabilization exercise, spinal manipulation, or traction treatment, providers may expect a high probability of a successful clinical outcome.

LBP imposes an enormous burden in the United States, both to individuals and to society. LBP is the most common type of pain reported by adults [1], and is among the most frequent complaints seen in physicians’ offices [2]. Moreover, 60% of LBP sufferers experience some form of functional limitation or disability as a result of their pain [3]. Pain and disability attributable to LBP are accompanied by an estimated \$100 billion to \$200 billion in health care expenditures and lost wages annually in the United States [4], equivalent to over 1% of the entire gross domestic product. Despite many recent advances in

*Corresponding author. Division of Physical Therapy, 520 Wakara Way, Salt Lake City, UT 84108. E-mail address: jeff.hebert@utah.edu (J. Hebert).

imaging and surgical technology, LBP prevalence and its related economic and societal burden have remained largely unchanged in the past decade [1,4].

Athletes may be especially susceptible to LBP and low back injuries. The prevalence of LBP appears particularly high for participants in sports that place high demands on the spine, such as wrestling, gymnastics, and golf [5]. Among the general population, LBP symptoms only weakly correlate with abnormal imaging findings and the great majority of cases of LBP cannot be attributed to specific pathoanatomical causes [6]. Athletes may be more likely than non-athletes to have an identifiable pathoanatomical cause of LBP symptoms [7,8]. Higher rates of spondylolysis, spondylolisthesis, and disc degeneration have been reported in athletes than in the general population [9,10]. Despite an increased incidence of certain pathoanatomical findings, it remains difficult to identify a specific cause in the majority of cases of LBP in athletes. The inability to identify a cause can make it difficult for clinicians to determine which treatment strategy is most likely to be effective. To assist clinicians in predicting which intervention is likely to be most effective, this article reviews the evidence for various interventions commonly used in the treatment of LBP.

SUB-GROUPING PATIENTS WITH LOW BACK PAIN

Common treatment alternatives for individuals with LBP, including those involved in athletics, consist of various forms of exercise, stabilization training, manual therapy, traction, and the use of physical modalities. Physical modalities, such as therapeutic ultrasound and electrical muscle stimulation, are widely used in the treatment of LBP [11], but randomized trials, systematic reviews, and practice guidelines have not supported the efficacy of these approaches [12–18]. Therefore, they are not considered as unique treatment strategies.

Exercise, manual therapy, traction, and many other treatments have been the subject of extensive scientific inquiry. Despite research efforts, evidence showing the effectiveness of these treatments is generally lacking or inconclusive. Even in studies that show some benefit for these treatments, the magnitude of the observed effects is often small, and the utility of these treatments remains subject to debate [13,16]. This can leave clinicians in a quandary as to the best treatment approach for a patient with LBP. The unfortunate result of this clinical dilemma is that one therapy can appear as appealing as the next, which may lead to less effective and efficient treatment. An increasing volume of information is available, however, to assist clinicians in predicting which type of treatment may be most likely to benefit an individual patient with LBP. Incorporation of this information into practice may improve clinical decision-making and treatment outcomes. A top priority for LBP research is to identify criteria for various subgroups of patients with LBP [19–21]. The nature of these subgroups and the methods for detecting them have been the subjects of a great deal of recent debate and research activity [22].

One approach to subgrouping patients with LBP has focused on identifying clusters of findings from the clinical examination that predict a more favorable outcome with a specific treatment approach [23]. Several experts in

rehabilitation, including McKenzie and others, have advocated this treatment-based approach. This article focuses on treatment-based subgrouping hypotheses originally described by Delitto and colleagues in 1995 [24]. The subgrouping hypotheses proposed are intended for patients who may or may not be involved in athletic activities with acute LBP or an acute exacerbation of LBP causing substantial pain and limitations in daily activities. After screening patients for any signs of serious pathology, information collected during the history and physical examination is used to place a patient into a subgroup. The name of each subgroup describes the fundamental treatment approach believed to offer the best chance for a successful outcome: manipulation, specific exercise (flexion, extension, and lateral shift patterns), stabilization, and traction. The cluster of examination findings and treatment strategies associated with each subgroup is reviewed in the following sections.

TREATMENT SUBGROUPS

Specific Exercise

The specific exercise subgroup emphasizes treatment using repeated end-range movements of the lumbar spine in a specific direction to affect the location and intensity of the patient's pain. This relationship between movement and pain was first emphasized by McKenzie [25]. Examination findings believed to identify patients in this subgroup include the presence of symptoms in the lower extremities, signs of nerve root compression (eg, positive straight-leg raise test; diminished reflex, sensation, or strength). The principle finding related to the specific exercise subgroup is the presence of centralization or a directional preference during the examination. Centralization occurs when a movement or position results in the relief of pain or paresthesia, or causes symptoms to move from a distal/lateral position in the buttocks and/or lower extremity to a more proximal location, closer to the midline of the lumbar spine [26]. Research has demonstrated the prognostic importance of the centralization phenomenon in patients with LBP with or without sciatica [27–33]. For example, Werneke and colleagues [32] examined the prognostic value of 23 demographic, psychosocial, occupational, and physical examination variables in 223 consecutive patients with acute LBP. The absence of centralization in this sample was associated with delayed recovery and the development of chronic LBP and disability. A concept related to centralization is directional preference. Directional preference occurs when a movement in one direction relieves pain or increases range of motion, and is often associated with movement in the opposite direction resulting in a worsening of the patient's signs and symptoms [34].

Advocates of a treatment-based classification approach contend that the presence of centralization or a directional preference is not just a favorable prognostic finding, but is among the predictive variables indicating the need for a specific exercise approach in treatment (Table 1). The basic treatment premise for patients in the specific exercise subgroup is the use of repeated, or sustained, end-range movements in the direction that caused centralization or of the directional preference determined during the examination. The movement

Table 1
Subgroups of patients with low back pain with subgroup criteria and treatment approaches

Subgroup	Subgroup criteria	Treatment approach
Specific exercise: extension	Symptoms distal to the buttock Symptoms centralize with lumbar extension Symptoms peripheralize with lumbar flexion Directional preference for extension	End-range extension exercises Mobilization to promote extension Avoidance of flexion activities
Specific exercise: flexion	Older age (>50 y) Directional preference for flexion Imaging evidence of lumbar spine stenosis	End-range flexion exercises Mobilization or manipulation of the spine and/or lower extremities Exercise to address impairments of strength or flexibility Body weight-supported ambulation
Stabilization	Younger age (<40 y) Average straight-leg raise (>91°) Aberrant movement present Positive prone-instability test	Exercises to strengthen large spinal muscles (erector spinae, oblique abdominals) Exercises to promote contraction of deep spinal muscles (multifidus, transversus abdominus)
Manipulation	No symptoms distal to knee Duration of symptoms <16 d Lumbar hypomobility Fear-Avoidance Beliefs Questionnaire for Work <19 Hip internal rotation range of motion >35°	Manipulation techniques for the lumbo-pelvic region Active lumbar range-of-motion exercises
Traction	Symptoms extend distal to the buttock(s) Signs of nerve root compression Peripheralization with extension movement; or positive contralateral straight-leg raise test	Prone mechanical traction Extension-specific exercises

may be flexion, extension, or lateral translation. The most common specific exercise movement for younger individuals or athletes is extension [35]. Treatments for patients who centralize or demonstrate a preference for extension include repeated end-range extension exercises, such as prone press-ups

(Fig. 1), or lumbar extension performed while standing. Exercises are progressed by increasing the amount of force or increasing the range of motion to maximize symptom relief. It is important that patients in this subgroup perform these activities frequently throughout the day. Patients may also need to be educated to avoid activities that promote prolonged or end-range flexion activities, such as lifting with poor body mechanics or sitting for long periods. Mobilization of the lumbar spine into extension (eg, posterior-to-anterior mobilization) may also be a useful treatment adjunct. An important contraindication to repeated end-range extension activities that should be considered in athletes with LBP is spondylolisthesis.

Several studies have investigated the effectiveness of an extension-specific exercise treatment approach. Studies that have applied this treatment to patients who fit the subgrouping criteria described above have reported evidence favoring the approach over other exercise interventions [36,37]. Trials that have evaluated an extension-specific exercise approach without an attempt to limit patients to those with these subgrouping criteria have generally not supported the effectiveness of this treatment option [38–41]. For example, Long and colleagues [35] randomized 230 patients with LBP who had a directional preference to receive either usual care, specific exercises in the direction of their preference, or specific exercises in the direction opposite their preference. The directional preference was extension for 83% of patients. Patients receiving exercises in the matched direction showed greater reductions in pain and disability after 2 weeks of treatment [35]. Browder and colleagues [35] randomized 48 patients with LBP who centralized with extension to receive either an extension-specific exercise approach or stabilization exercises. Patients receiving the extension-specific exercise approach, which included extension exercises, patient education, and graded posterior-to-anterior mobilization, showed greater improvement in disability at both short- (1- and 4-week), and long-term (6-month) outcomes [35].

Stabilization Exercise

Functional deficits of the trunk muscles have been observed in general [42,43,44] and athletic [45,46] populations with LBP. Similar deficits in trunk



Fig. 1. Example of specific exercise: prone press-up.

muscle function have also been associated with traumatic knee injury [47,48] and chronic groin pain in athletes [49], suggesting that a lack of trunk control may compromise function or stability of the lower extremities during athletic activity. In addition to deficits in neuromuscular control of the trunk muscles, patients with LBP have also been observed to have morphologic changes, including atrophy [50–54] and fatty infiltration [55–57] in the lumbar multifidus and erector spinae muscles. Stabilization exercise programs are typically designed to address the deficits in strength, endurance, and function of the trunk musculature that have been identified in patients with LBP. It is thought that improvements in trunk muscle function lead to decreases in pain and disability by improving the control of spinal segments during movement. In support of this hypothesis, stabilization exercise has been shown to improve trunk muscle function [58,59] and morphology [60–62] in individuals with LBP.

Improvements in trunk muscle function and morphology may represent important outcomes of rehabilitation programs. However, these physiologic changes may not correspond to patient-centered improvements in pain and disability. This concern is highlighted by the conflicting results of research examining the effects of stabilization exercise programs on patient-centered outcomes. While some studies support stabilization exercises as an effective treatment for LBP [63–66], others have demonstrated equivalence between stabilization exercise and traditional rehabilitation approaches [67] or manual therapy [68,69]. A recent systematic review by Rackwitz and colleagues [70] concluded that stabilization exercise for LBP is more effective than treatment by a general practitioner but not more effective than other physiotherapy interventions.

Conflicting findings of research evaluating the effectiveness of stabilization exercise support the consideration that there may be a subgroup of patients with LBP who are most likely to benefit from this approach. Hicks and colleagues [71] investigated variables that may identify which patients with LBP are likely to experience clinical success when receiving stabilization exercises. Four variables (see Table 1) were most predictive of success, defined as a 50% reduction in disability as measured by the Oswestry Questionnaire. When three or more of these variables were present, the probability of achieving clinical success increased from 33% to 67%. While the presence of these variables was associated with an increased likelihood of success, it is clear that future research may be able to identify additional factors to improve the prediction of success with stabilization exercises.

The most effective exercises for use in a stabilization program are also a matter of current debate. A good deal of recent attention and research has focused on specific retraining exercises for the deep trunk muscles, in particular the transversus abdominus and multifidus [58,60,64]. The goal of this approach is to retrain the normal stabilizing motor patterns of these muscles, which are often compromised in individuals with LBP. Some evidence for this approach exists [58,64]. However, most studies have compared this specific retraining approach to management involving no exercise or poorly defined

exercise protocols. Other stabilization regimens have placed greater emphasis on exercises designed to improve the strength and endurance of larger, more superficial trunk muscles (ie, erector spinae, oblique abdominals, quadratus lumborum) (Fig. 2) [70,71]. This approach to stabilization exercise emphasizes the use of strengthening exercises that sufficiently challenge these important muscle groups while minimizing potentially harmful compressive and shear loading of the spine. Since the stabilizing activity of any of these muscles is generally that of a low-intensity contraction [72], exercise protocols focus on high repetitions of low-load contractions to promote muscle endurance. Recent research has compared specific muscle retraining programs to this more general stabilization approach [61,63,67]. These studies have not found differences favoring one approach over the other. Although many experts advocate the necessity of specifically retraining the deep spinal muscles, the evidence does not clearly support this perspective. Further research should help define the optimal mix of stabilization exercises for patients with LBP.

Spinal Manipulation

Spinal manipulation is generally defined as the application of a high-velocity, low-amplitude thrust to a joint, which frequently results in an audible “crack” or cavitation [73]. The clinical outcomes associated with spinal manipulation have been the subjects of a great deal of scientific investigation. This research has resulted in several randomized clinical trials [74–77] and systematic reviews [16,78] demonstrating effectiveness for manipulation when compared with placebo or other interventions. However, other trials and reviews have failed to



Fig. 2. Examples of stabilization exercises: side bridge (*above*) and bird dog (*below*).

demonstrate a clear clinical benefit of spinal manipulation when compared with other therapies [37,79–81]. Clinical experience suggests that spinal manipulation is effective for at least some patients with LBP. The conflict between research outcomes and clinical experience may be due in part to uncertainty in defining which subgroup of patients with LBP is most likely to benefit from manipulation [82].

Manipulation has been used for centuries [83], yet the mechanism by which manipulation may have a therapeutic effect is subject to debate, which leads to confusion in determining the subgroup that responds best to the treatment. Traditional explanations of the therapeutic mechanism of spinal manipulation have emphasized the importance of directing forces to specific spinal joints for the purpose of correcting a biomechanical dysfunction or misalignment [84–87]. While these constructs may seem intuitive, several studies have questioned their validity. Research has questioned the ability of clinicians to direct manipulative forces in a manner to affect specific spinal joints [88,89]. Furthermore, while there is a small degree of intervertebral movement produced during spinal manipulation [90–92], sustained changes in alignment have not been observed [93].

Flynn and colleagues [94] used a different approach to examining the patient characteristics that may define a subgroup of patients likely to benefit from manipulation by focusing on the prediction of clinical success instead of presumptions based on biomechanical theories. This study identified five variables (see Table 1) predictive of success, defined as a 50% reduction in the Oswestry Questionnaire within 1 week. Patients were considered to be likely responders to manipulation when four or more of these variables were present. When patients met this threshold, the probability of achieving clinical success increased from 45% to 95%.

A follow-up study [76] was performed to examine the validity of these predictive criteria. The results found that patients with LBP receiving manipulation who met these criteria experienced greater decreases in pain and disability than did patients who received manipulation but did not meet the criteria. Additionally, patients who met the criteria and received manipulation experienced greater improvement than did patients who met the criteria but were treated with stabilization exercises. Two important conclusions can be inferred from this study. First, while results do not support spinal manipulation as a superior treatment for all patients with LBP, they do suggest that manipulation is effective for the appropriate subgroup of patients. Second, the presence of these criteria does not automatically equate to a more favorable prognosis unless the appropriate treatment is provided.

The manipulative technique used in this research (Fig. 3) involves a high-velocity, low-amplitude thrust delivered to the anterior superior iliac spine of the supine patient after being sidebent away from and rotated toward the clinician. This procedure was originally thought to be appropriate for patients with biomechanical dysfunction of the sacroiliac articulation [24]. Except for the presence of stiffness somewhere in the lumbar spine, none of the variables



Fig. 3. Manipulation technique.

predictive of success with this procedure related to biomechanical dysfunction. The lack of relationship between clinical success and specific biomechanical dysfunction seems consistent with the growing body of evidence supporting a primarily neurophysiologic mechanism of manipulation. Studies have shown spinal manipulation to affect both sensory and motor nerve activity as well as electromyographic-measured muscle activity [92,95–101]. Although further research is needed, one conclusion can be drawn: If the mechanism of action for spinal manipulation is primarily mediated by neurophysiologic mechanisms rather than biomechanical “realignment,” clinicians may be forced to change their paradigm for determining which patients are most appropriate for spinal manipulation.

Traction

Traction techniques for the lumbar spine have a rich history in medicine dating back more than 200 years [102]. Lumbar traction is commonly used [103] and has been referred to by some as “decompression therapy” [104]. Many clinicians believe traction is effective [103,105,106], but the usefulness of traction for treating LBP has been the subject of debate and controversy [107,108].

The traditional presumption of clinicians has considered the presence of sciatica or signs of nerve root compression as indications for traction [103]. Yet, until recently, little research evidence has been available to assist clinicians in predicting which patients with LBP were most likely to benefit from traction. We recently examined the outcomes of patients with LBP who also had symptoms below the buttock and signs of nerve root compression. Our purpose was to determine if these criteria were specific enough to define the subgroup of patients who respond to traction, or if additional criteria were required [27]. We found that the presence of symptoms below the buttock and signs of nerve root compression were not specific enough to define this subgroup of patients. Two additional factors were found to identify patients likely to respond favorably to

traction: (1) peripheralization with extension movement and (2) a positive crossed (ie, contralateral) straight-leg raise test. The presence of either of these, in addition to symptoms below the buttock and signs of nerve root compression, define the subgroup of patients who respond to traction. Peripheralization occurs when a movement or posture causes symptoms to move distally, away from the spinal midline. A positive crossed straight-leg raise test is defined as reproduction of the patient's familiar lower extremity symptoms when the contralateral leg is passively raised with the knee maintained in an extended position [109]. When patients with symptoms below the buttock and signs of nerve root compression had either of these findings and received traction along with an extension-specific exercise program, they showed greater short-term reductions in disability than patients with these findings who received only the extension exercise program. These results suggest that traction may be essential to maximize improvement in a specific subgroup of patients (see [Table 1](#)).

There has been considerable diversity in the recommended parameters to be used when applying traction. The most common patient position used with mechanical traction is reported to be supine with the hips and knees flexed approximately 90° [103,110]. Although this position is comfortable for many patients with LBP, the position places the lumbar spine in flexion and may therefore be contraindicated for patients who meet the traction subgroup criteria. Prone lying ([Fig. 4](#)) may therefore be a preferred position for these patients. There is no clear evidence regarding the most effective traction force. Many experts contend that the force needs to be higher than is typically used in clinical practice (~50% of body weight) to produce a therapeutic effect [111,112]. It may be appropriate to initiate treatment at a slightly lower force (~40% of body weight), then increase the force as tolerated up to a maximum of 60% of body weight. With high-force traction, the duration of treatment may need



Fig. 4. Traction therapy.

to be shorter (8–12 minutes), with allowances for ramping up and ramping down the force. Because the goal of traction is vertebral separation, static traction is often recommended [111,112]. Traction is rarely delivered as a stand-alone treatment. Because the overall goals of treatment for patients in this subgroup are to reduce and centralize leg symptoms, traction is frequently delivered along with an extension-specific exercise program as described.

Evidence-based guidelines and systematic reviews have not supported the effectiveness of traction for patients with LBP [16,113,114]. The discrepancy between clinical perceptions and research evidence may be attributable to the manner in which traction has been applied in the majority of studies that have examined its effectiveness. Studies that have shown no benefit from using traction have used nonspecific inclusion criteria, essentially allowing all patients fitting a broad definition of acute or chronic LBP to enter [115,116]. Most studies have also failed to adequately define the parameters used for delivering the traction, or have used parameters that are not consistent with expert opinions or typical clinical use [108]. To use traction most effectively, greater attention is needed in the identification of clinical factors that pinpoint patients who need traction and in the application of appropriate dosages of traction.

SUMMARY

The identification of predictive factors in patients with LBP should allow the patient to be matched with the most appropriate treatment intervention, maximizing the likelihood of a favorable clinical outcome [117]. While the identification of predictive factors for the treatment of patients with LBP represents a significant advance in patient care, much more information and research are needed. Nevertheless, it appears that using simple baseline evaluation findings can help clinicians more efficiently and effectively select the most appropriate treatment for an individual patient with LBP.

References

- [1] Deyo RA, Mirza SK, Martin BI. Back pain prevalence and visit rates: estimates from U.S. national surveys, 2002. *Spine* 2006;31(23):2724–7.
- [2] Hart LG, Deyo RA, Cherkin DC. Physician office visits for low back pain. Frequency, clinical evaluation, and treatment patterns from a U.S. national survey. *Spine* 1995;20(1):11–9.
- [3] Heliövaara M, Sievers K, Impivaara O, et al. Descriptive epidemiology and public health aspects of low back pain. *Ann Med* 1989;21(5):327–33.
- [4] Katz JN. Lumbar disc disorders and low-back pain: socioeconomic factors and consequences. *J Bone Joint Surg Am* 2006;88(Suppl 2):21–4.
- [5] Bono CM. Low-back pain in athletes. *J Bone Joint Surg Am* 2004;86(2):382–96.
- [6] van Tulder MW, Assendelft WJ, Koes BW, et al. Spinal radiographic findings and nonspecific low back pain. A systematic review of observational studies. *Spine* 1997;22(4):427–34.
- [7] Hardcastle P, Annear P, Foster DH, et al. Spinal abnormalities in young fast bowlers. *J Bone Joint Surg Br* 1992;74(3):421–5.
- [8] Sward L, Hellstrom M, Jacobsson B, et al. Disc degeneration and associated abnormalities of the spine in elite gymnasts. A magnetic resonance imaging study. *Spine* 1991;16(4):437–43.

- [9] Ong A, Anderson J, Roche J. A pilot study of the prevalence of lumbar disc degeneration in elite athletes with lower back pain at the Sydney 2000 Olympic Games. *Br J Sports Med* 2003;37(3):263–6.
- [10] McCarroll JR, Miller JM, Ritter MA. Lumbar spondylolysis and spondylolisthesis in college football players. A prospective study. *Am J Sports Med* 1986;14(5):404–6.
- [11] Wong RA, Schumann B, Townsend R, et al. A survey of therapeutic ultrasound use by physical therapists who are orthopaedic certified specialists. *Phys Ther* 2007;87(8):986–94 [discussion: 995–1001].
- [12] Khadilkar A, Milne S, Brosseau L, et al. Transcutaneous electrical nerve stimulation for the treatment of chronic low back pain: a systematic review. *Spine* 2005;30(23):2657–66.
- [13] van Tulder MW, Koes B, Malmivaara A. Outcome of non-invasive treatment modalities on back pain: an evidence-based review. *Eur Spine J* 2006;15(Suppl 1):S64–81.
- [14] Khadilkar A, Milne S, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic low-back pain. *Cochrane Database Syst Rev* 2005;(3):CD003008.
- [15] Hurwitz EL, Morgenstern H, Kominski GF, et al. A randomized trial of chiropractic and medical care for patients with low back pain: eighteen-month follow-up outcomes from the UCLA low back pain study. *Spine* 2006;31(6):611–21 [discussion: 622].
- [16] Chou R, Huffman LH. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med* 2007;147(7):492–504.
- [17] Philadelphia Panel. Philadelphia Panel evidence-based clinical practice guidelines on selected rehabilitation interventions for low back pain. *Phys Ther* 2001;81(10):1641–74.
- [18] Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med* 2007;147(7):478–91.
- [19] Borkan JM, Cherkin DC. An agenda for primary care research on low back pain. *Spine* 1996;21(24):2880–4.
- [20] Borkan JM, Koes B, Reis S, et al. A report from the second international forum for primary care research on low back pain: reexamining priorities. *Spine* 1998;23(18):1992–6.
- [21] Henschke N, Maher CG, Refshauge KM, et al. Low back pain research priorities: a survey of primary care practitioners. *BMC Fam Pract* 2007;8:40.
- [22] Billis EV, McCarthy CJ, Oldham JA. Subclassification of low back pain: a cross-country comparison. *Eur Spine J* 2007;16(7):865–79.
- [23] Riddle DL. Classification and low back pain: a review of the literature and critical analysis of selected systems. *Phys Ther* 1998;78(7):708–37.
- [24] Delitto A, Erhard RE, Bowling RW. A treatment-based classification approach to low back syndrome: identifying and staging patients for conservative management. *Phys Ther* 1995;75:470–89.
- [25] McKenzie RA. The lumbar spine: mechanical diagnosis and therapy. Waikanae (New Zealand): Spinal Publications Limited; 1989.
- [26] Fritz JM, Cleland JA, Childs JD. Subgrouping patients with low back pain: evolution of a classification approach to physical therapy. *J Orthop Sports Phys Ther* 2007;37(6):290–302.
- [27] Fritz JM, Lindsay W, Matheson JW, et al. Is there a subgroup of patients with low back pain likely to benefit from mechanical traction? Results of a randomized clinical trial and subgrouping analysis. *Spine* 2007;32(26):E793–800.
- [28] Berthelot JM, Delecrin J, Maugars Y, et al. Contribution of centralization phenomenon to the diagnosis, prognosis, and treatment of diskogenic low back pain. *Joint Bone Spine* 2007;74(4):319–23.
- [29] George SZ, Bialosky JE, Donald DA. The centralization phenomenon and fear-avoidance beliefs as prognostic factors for acute low back pain: a preliminary investigation involving patients classified for specific exercise. *J Orthop Sports Phys Ther* 2005;35(9):580–8.

- [30] Karas R, McIntosh G, Hall H, et al. The relationship between nonorganic signs and centralization of symptoms in the prediction of return to work for patients with low back pain. *Phys Ther* 1997;77:354–60.
- [31] Long AL. The centralization phenomenon. Its usefulness as a predictor or outcome in conservative treatment of chronic low back pain (a pilot study). *Spine* 1995;20(23):2513–20 [discussion: 2521].
- [32] Werneke M, Hart DL. Centralization phenomenon as a prognostic factor for chronic low back pain and disability. *Spine* 2001;26(7):758–64 [discussion: 765].
- [33] Skytte L, May S, Petersen P. Centralization: its prognostic value in patients with referred symptoms and sciatica. *Spine* 2005;30(11):E293–9.
- [34] Kilpikoski S, Airaksinen O, Kankaanpää M, et al. Interexaminer reliability of low back pain assessment using the McKenzie method. *Spine* 2002;27:E207–14.
- [35] Long AL, Donelson R. Does it matter which exercise? A randomized trial of exercise for low back pain. *Spine* 2004;29:2593–602.
- [36] Browder DA, Childs JD, Cleland JA, et al. Effectiveness of an extension-oriented treatment approach in a subgroup of subjects with low back pain: a randomized clinical trial. *Phys Ther* Sep 2007;87(12):1608–18.
- [37] Machado LA, de Souza MS, Ferreira PH, et al. The McKenzie method for low back pain: a systematic review of the literature with a meta-analysis approach. *Spine* 2006;31(9):E254–62.
- [38] Cherkin DC, Deyo RA, Battie M, et al. A comparison of physical therapy, chiropractic manipulation, and provision of an educational booklet for the treatment of patients with low back pain. *N Engl J Med* 1998;339(15):1021–9.
- [39] Dettori JR, Bullock SH, Suttlive TG, et al. The effects of spinal flexion and extension exercises and their associated postures in patients with acute low back pain. *Spine* 1995;20(21):2303–12.
- [40] Indahl A, Velund L, Reikeraas O. Good prognosis for low back pain when left untampered. A randomized clinical trial. *Spine* 1995;20(4):473–7.
- [41] Malmivaara A, Hakkinen U, Aro T, et al. The treatment of acute low back pain—bed rest, exercises, or ordinary activity? *N Engl J Med* 1995;332(6):351–5.
- [42] Hodges PW, Richardson CA. Contraction of the abdominal muscles associated with movement of the lower limb. *Phys Ther* 1997;77(2):132–42.
- [43] Hodges PW, Richardson CA. Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine* 1996;21:2640–50.
- [44] Newcomer KL, Laskowski ER, Yu B, et al. Differences in repositioning error among patients with low back pain compared with control subjects. *Spine* 2000;25(19):2488–93.
- [45] Reeves NP, Cholewicki J, Silfies SP. Muscle activation imbalance and low-back injury in varsity athletes. *J Electromyogr Kinesiol* 2006;16(3):264–72.
- [46] Renkawitz T, Boluki D, Grifka J. The association of low back pain, neuromuscular imbalance, and trunk extension strength in athletes. *Spine J* 2006;6(6):673–83.
- [47] Zazulak BT, Hewett TE, Reeves NP, et al. Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *Am J Sports Med* 2007;35(7):1123–30.
- [48] Zazulak BT, Hewett TE, Reeves NP, et al. The effects of core proprioception on knee injury: a prospective biomechanical-epidemiological study. *Am J Sports Med* 2007;35(3):368–73.
- [49] Cowan SM, Schache AG, Brukner P, et al. Delayed onset of transversus abdominis in long-standing groin pain. *Med Sci Sports Exerc* 2004;36(12):2040–5.
- [50] Barker KL, Shamley DR, Jackson D. Changes in the cross-sectional area of multifidus and psoas in patients with unilateral back pain: the relationship to pain and disability. *Spine* 2004;29(22):E515–9.

- [51] Danneels LA, Vanderstraeten GG, Cambier DC, et al. CT imaging of trunk muscles in chronic low back pain patients and healthy control subjects. *Eur Spine J* 2000;9(4): 266–72.
- [52] Hides JA, Stokes MJ, Saide M, et al. Evidence of lumbar multifidus muscle wasting ipsilateral to symptoms in patients with acute/subacute low back pain. *Spine* 1994;19(2): 165–72.
- [53] Ng JK, Richardson CA, Kippers V, et al. Relationship between muscle fiber composition and functional capacity of back muscles in healthy subjects and patients with back pain. *J Orthop Sports Phys Ther* 1998;27(6):389–402.
- [54] Yoshihara K, Nakayama Y, Fujii N, et al. Atrophy of the multifidus muscle in patients with lumbar disk herniation: histochemical and electromyographic study. *Orthopedics* 2003;26(5):493–5.
- [55] Kang CH, Shin MJ, Kim SM, et al. MRI of paraspinal muscles in lumbar degenerative kyphosis patients and control patients with chronic low back pain. *Clin Radiol* 2007;62(5):479–86.
- [56] Kjaer P, Bendix T, Sorensen JS, et al. Are MRI-defined fat infiltrations in the multifidus muscles associated with low back pain? *BMC Med* 2007;5:2.
- [57] Mengiardi B, Schmid MR, Boos N, et al. Fat content of lumbar paraspinal muscles in patients with chronic low back pain and in asymptomatic volunteers: quantification with MR spectroscopy. *Radiology* 2006;240(3):786–92.
- [58] Tsao H, Hodges PW. Persistence of improvements in postural strategies following motor control training in people with recurrent low back pain. *J Electromyogr Kinesiol* 2007, in press.
- [59] Tsao H, Hodges PW. Immediate changes in feedforward postural adjustments following voluntary motor training. *Exp Brain Res* 2007;181(4):537–46.
- [60] Hides JA, Richardson CA, Jull GA. Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine* 1996;21(23):2763–9.
- [61] Danneels LA, Vanderstraeten GG, Cambier DC, et al. Effects of three different training modalities on the cross sectional area of the lumbar multifidus muscle in patients with chronic low back pain. *Br J Sports Med* 2001;35(3):186–91.
- [62] Rissanen A, Kalimo H, Alaranta H. Effect of intensive training on the isokinetic strength and structure of lumbar muscles in patients with chronic low back pain. *Spine* 1995;20(3): 333–40.
- [63] Koumantakis GA, Watson PJ, Oldham JA. Trunk muscle stabilization training plus general exercise versus general exercise only: randomized controlled trial of patients with recurrent low back pain. *Phys Ther* 2005;85(3):209–25.
- [64] Koumantakis GA, Watson PJ, Oldham JA. Supplementation of general endurance exercise with stabilisation training versus general exercise only. Physiological and functional outcomes of a randomised controlled trial of patients with recurrent low back pain. *Clin Biomech (Bristol, Avon)* 2005;20(5):474–82.
- [65] Hides JA, Jull GA, Richardson CA. Long-term effects of specific stabilizing exercises for first-episode low back pain. *Spine* 2001;26(11):E243–8.
- [66] O'Sullivan PB, Phytz GD, Twomey LT, et al. Evaluation of specific stabilizing exercise in the treatment of chronic low back pain with radiologic diagnosis of spondylolysis or spondylolisthesis. *Spine* 1997;22(24):2959–67.
- [67] Cairns MC, Foster NE, Wright C. Randomized controlled trial of specific spinal stabilization exercises and conventional physiotherapy for recurrent low back pain. *Spine* 2006;31(19):E670–81.
- [68] Goldby LJ, Moore AP, Doust J, et al. A randomized controlled trial investigating the efficiency of musculoskeletal physiotherapy on chronic low back disorder. *Spine* 2006;31(10):1083–93.
- [69] Rasmussen-Barr E, Nilsson-Wikmar L, Arvidsson I. Stabilizing training compared with manual treatment in sub-acute and chronic low-back pain. *Man Ther* 2003;8(4):233–41.

- [70] Rackwitz B, de Bie R, Limm H, et al. Segmental stabilizing exercises and low back pain. What is the evidence? A systematic review of randomized controlled trials. *Clin Rehabil* 2006;20(7):553–67.
- [71] Hicks GE, Fritz JM, Delitto A, et al. Preliminary development of a clinical prediction rule for determining which patients with low back pain will respond to a stabilization exercise program. *Arch Phys Med Rehabil* 2005;86(9):1753–62.
- [72] McGill SM. Low back exercises: evidence for improving exercise regimens. *Phys Ther* 1998;78(7):754–65.
- [73] Evans DW. Mechanisms and effects of spinal high-velocity, low-amplitude thrust manipulation: previous theories. *J Manipulative Physiol Ther* 2002;25(4):251–62.
- [74] Aure OF, Nilsen JH, Vasseljen O. Manual therapy and exercise therapy in patients with chronic low back pain: a randomized, controlled trial with 1-year follow-up. *Spine* 2003;28(6):525–31 [discussion: 531–2].
- [75] Team UBT. United Kingdom back pain exercise and manipulation (UK BEAM) randomised trial: effectiveness of physical treatments for back pain in primary care. *BMJ* 2004;329:1377–85.
- [76] Childs JD, Fritz JM, Flynn TW, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med* 2004;141(12):920–8.
- [77] Giles LG, Muller R. Chronic spinal pain: a randomized clinical trial comparing medication, acupuncture, and spinal manipulation. *Spine* 2003;28(14):1490–502.
- [78] Assendelft WJ, Koes BW, van der Heijden GJ, et al. The efficacy of chiropractic manipulation for back pain: blinded review of relevant randomized clinical trials. *J Manipulative Physiol Ther* 1992;15(8):487–94.
- [79] Assendelft WJ, Morton SC, Yu EI, et al. Spinal manipulative therapy for low back pain. *Cochrane Database Syst Rev* 2004;(1):CD000447.
- [80] Cherkin DC, Sherman KJ, Deyo RA, et al. A review of the evidence for the effectiveness, safety, and cost of acupuncture, massage therapy, and spinal manipulation for back pain. *Ann Intern Med* 2003;138(11):898–906.
- [81] Hurwitz EL, Morgenstern H, Harber P, et al. A randomized trial of medical care with and without physical therapy and chiropractic care with and without physical modalities for patients with low back pain: 6-month follow-up outcomes from the UCLA low back pain study. *Spine* 2002;27(20):2193–204.
- [82] Bouter LM, van Tulder MW, Koes BW. Methodologic issues in low back pain research in primary care. *Spine* 1998;23:2014–20.
- [83] Curtis P. Spinal manipulation: does it work? *Occup Med* 1988;3(1):31–44.
- [84] Bernard H. The mechanism of anatomical structure in its relation to osteopathy. 1911. *J Am Osteopath Assoc* 2000;100(7):444–8.
- [85] Cyriax JH. Diagnosis of soft tissue lesions. In: Cyriax JH, editor. *Textbook of orthopaedic medicine*. 6th edition. Baltimore (MD): Williams & Wilkins; 1976. p. 389.
- [86] Maitland GD. *Vertebral manipulation*. 5th edition. Oxford (UK): Butterworth Heinemann; 1986.
- [87] Meeker WC, Haldeman S. Chiropractic: a profession at the crossroads of mainstream and alternative medicine. *Ann Intern Med* 2002;136(3):216–27.
- [88] Ross JK, Bereznick DE, McGill SM. Determining cavitation location during lumbar and thoracic spinal manipulation: Is spinal manipulation accurate and specific? *Spine* 2004;29(13):1452–7.
- [89] Bereznick DE, Ross JK, McGill SM. The frictional properties at the thoracic skin-fascia interface: implications in spine manipulation. *Clin Biomech (Bristol, Avon)* 2002;17(4):297–303.
- [90] Gal J, Herzog W, Kawchuk G, et al. Movements of vertebrae during manipulative thrusts to unembalmed human cadavers. *J Manipulative Physiol Ther* 1997;20(1):30–40.

- [91] Maigne JY, Guillon F. Highlighting of intervertebral movements and variations of intradiskal pressure during lumbar spine manipulation: a feasibility study. *J Manipulative Physiol Ther* 2000;23:531–5.
- [92] Colloca CJ, Keller TS, Gunzburg R. Biomechanical and neurophysiological responses to spinal manipulation in patients with lumbar radiculopathy. *J Manipulative Physiol Ther* 2004;27(1):1–15.
- [93] Tullberg T, Blomberg S, Branth B, et al. Manipulation does not alter the position of the sacroiliac joint. A roentgen stereophotogrammetric analysis. *Spine* 1998;23(10):1124–8.
- [94] Flynn T, Fritz J, Whitman J, et al. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with spinal manipulation. *Spine* 2002;27(24):2835–43.
- [95] Colloca CJ, Keller TS, Gunzburg R. Neuromechanical characterization of in vivo lumbar spinal manipulation. Part II. Neurophysiological response. *J Manipulative Physiol Ther* 2003;26(9):579–91.
- [96] Dishman JD, Ball KA, Burke J. First prize: central motor excitability changes after spinal manipulation: a transcranial magnetic stimulation study. *J Manipulative Physiol Ther* 2002;25(1):1–9.
- [97] Dishman JD, Cunningham BM, Burke J. Comparison of tibial nerve H-reflex excitability after cervical and lumbar spine manipulation. *J Manipulative Physiol Ther* 2002;25(5):318–25.
- [98] Herzog W, Scheele D, Conway PJ. Electromyographic responses of back and limb muscles associated with spinal manipulative therapy. *Spine* 1999;24(2):146–52 [discussion: 153].
- [99] Suter E, McMorland G, Herzog W, et al. Decrease in quadriceps inhibition after sacroiliac joint manipulation in patients with anterior knee pain. *J Manipulative Physiol Ther* 1999;22(3):149–53.
- [100] Suter E, McMorland G, Herzog W, et al. Conservative lower back treatment reduces inhibition in knee-extensor muscles: a randomized controlled trial. *J Manipulative Physiol Ther* 2000;23(2):76–80.
- [101] Suter E, McMorland G, Herzog W. Short-term effects of spinal manipulation on H-reflex amplitude in healthy and symptomatic subjects. *J Manipulative Physiol Ther* 2005;28(9):667–72.
- [102] Silver J. The history of modern spinal traction with particular reference to neural disorders. *Spinal Cord* 1997;35(10):710–1.
- [103] Harte AA, Gracey JH, Baxter GD. Current use of lumbar traction in the management of low back pain: results of a survey of physiotherapists in the United Kingdom. *Arch Phys Med Rehabil* 2005;86(6):1164–9.
- [104] Daniel DM. Non-surgical spinal decompression therapy: Does the scientific literature support efficacy claims made in the advertising media? *Chiropr Osteopat* 2007;15:7.
- [105] Mikhail C, Korner-Bitensky N, Rossignol M, et al. Physical therapists' use of interventions with high evidence of effectiveness in the management of a hypothetical typical patient with acute low back pain. *Phys Ther* 2005;85:1151–67.
- [106] Poitras S, Blais R, Swaine B, et al. Management of work-related low back pain: a population-based survey of physical therapists. *Phys Ther* 2005;85:1168–81.
- [107] Clarke JA, van Tulder MW, Blomberg SE, et al. Traction for low-back pain with or without sciatica. *Cochrane Database Syst Rev* 2007;(2):CD003010.
- [108] Harte AA, Baxter GD, Gracey JH. The efficacy of traction for back pain: a systematic review of randomized controlled trials. *Arch Phys Med Rehabil* 2003;84(10):1542–53.
- [109] Kosteljanetz M, Bang F, Schmidt-Olsen S. The clinical significance of straight-leg raising (Lasegue's sign) in the diagnosis of prolapsed lumbar disc. Interobserver variation and correlation with surgical finding. *Spine* 1988;13(4):393–5.

- [110] Harte AA, Baxter GD, Gracey JH. The effectiveness of motorized lumbar traction in the management of LBP with lumbo sacral nerve root involvement: a feasibility study. *BMC Musculoskelet Disord* 2007;8:118.
- [111] Judovich B. Lumbar traction therapy. *JAMA* 1955;159:549–52.
- [112] Saunders HD. Evaluation, treatment and prevention of musculoskeletal disorders, Vol. 1. 4th edition. Chaska (MN): The Saunders Group; 2004.
- [113] Clarke J, van Tulder M, Blomberg S, et al. Traction for low back pain with or without sciatica: an updated systematic review within the framework of the Cochrane collaboration. *Spine* 2006;31:1591–9.
- [114] Macario A, Pergolizzi JV. Systematic literature review of spinal decompression via motorized traction for chronic discogenic low back pain. *Pain Pract* 2006;6:171–8.
- [115] Beurskens AJ, De Vet HC, Koke AJ, et al. Efficacy of traction for nonspecific low back pain. 12-week and 6-month results of a randomized clinical trial. *Spine* 1997;22:2756–62.
- [116] Werners R, Pynsent PB, Bulstrode CJK. Randomized trial comparing interferential therapy with motorized lumbar traction and massage in the management of low back pain in a primary care setting. *Spine* 1999;24:1579–84.
- [117] Brennan GP, Fritz JM, Hunter SJ, et al. Identifying subgroups of patients with acute/subacute “nonspecific” low back pain: results of a randomized clinical trial. *Spine* 2006;31(6):623–31.